PELAGIC COPEPODS FROM THE INSHORE WATERS OFF SAURASHTRA COAST

M. I. PATEL

Marine biological Research Station, Port Okha, India

ABSTRACT

Pelagic copepods from the coastal waters of Saurashtra are studied. Seventy eight species of copepods belonging to the suborder Calanoida, Harpacticoida and Cyclopoida are identified. The Calanoida is represented by 15 families, 23 genera and 54 species; the Harpacticoida by 6 families, 7 genera and 7 species and Cyclopoida by 4 families, 5 genera and 17 species. *Pontella karachiensis* Fazal-ur-Rehman is reported here for the first time from the Indian Coast.

INTRODUCTION

PELAGIC COPEPODS from the inshore waters have been fairly well investigated from the east and southwest coasts of India (Prasad, 1964), whereas these from the northwest coast are inadequately known. A preliminary account of the plankton off Kandla Port (Lat. 23° 01' N, 70° 13' E) in the Gulf of Kutch was given by Ramamurthy and Dhawan (1963). Recently Bhaskaran and Gopalakrishnan (1971) made a preliminary study on the marine plankton of the Gulf of Kutch, off Port Okha (Lat. 22° 28' N, 69° 05' E).

The author has made an attempt to give a systematic list of the pelagic copepods inhabiting the coastal waters of Saurashtra. The classification adopted is that of Sars (1903, 1911, 1918). The present study was carried out as a part of Plankton Project, between 1972 and 1974 by the author at the Marine Biological Research Station, Port Okha.

The author is grateful to Shri Moosa Raza, I.A.S., Commissioner of Fisheries, Government of Gujarat for facilities provided. He is thankful to Shri M. Bhaskaran, Research Officer for encouragement. Thanks are also due to Shri I. N. Mahyavanshi for offering plankton samples collected by him from November 1972 to July 1973.

AREA OF STUDY

The study area extends between the Latitudes $20^{\circ}50' 28''N$ and $22^{\circ}30' 18''N$ and Longitudes $68^{\circ}50'E$ and $70^{\circ}20'10''E$, covering a stretch of 360 km of the coast. Stations in the neritic waters of the Saurashtra Coast from where samples have been collected are given in Table 1.

Sampling station	Location	Approximate distance from shore (in nautical miles)	Approximate depth (m)	
Off Rozi	22-70 4A	4.3	6-14	
Off Port Okha	22-69 3A	1.6	6	
Off Dwarka	22-68 2E, 2F	6	38-46	
Off Porbandar	21-69 3C, 4C	10	34-46	
Off Veraval	20-70 5A, 6A	15	36-48	

TABLE 1. Location of stations

MATERIALS AND METHODS

The collections were made by an half metre organdy cloth net having 36 Strands/ cm, towed horizontally for 20 minutes duration in the surface waters from a survey cum exploratory vessel or a 'machhawa' with out-board motor. In all, seventy six samples were collected. All plankton samples were preserved in 5% formaldehyde and examined for species composition. Sketches were drawn with the aid of a camera lucida.

LIST OF SPECIES

SUBORDER: CALANOIDA

TRIBE: AMPHASCANDRIA

Family: CALANIDAE

Genus: Canthocalanus A. Scott Canthocalanus pauper (Giesbrecht)

Genus: Undinula A. Scott Undinula vulgaris (Dana)

(a) variety typica Sewell
 (b) variety giesvrechti Sewell

Undinula darwini (Lubbock) (a) variety stmmetrica Sewell (b) variety intermedia Sewell

Family: EUCALANIDAE

Genus: Eucalanus Dana

Eucalanus elongatus (Dana)

E. monachus Giesbrecht

- E. crassus Giesbrecht
- E. subcrassus Giesbrecht E. attenuatus (Dana)
- E. pseudattenuatus (Dana)
- Genus: Rhincalanus Dana Rhincalanus cornutus Dana R. nasutus Giesbrecht

Family: PARACALANIDAE

Genus: Paracalanus Boeck Paracalanus parvus Giesbrecht P. aculeatus Giesbrecht

P. denudatus Sewell

Genus: Acrocalanus Giesbrecht Acrocalanus gracilis Giesbrecht A. gibber Giesbrecht A. longicornis Giesbrecht

Family: PSEUDOCALANIDAE

Genus: Calocalanus Giesbrecht Calocalanus pavo (Dana) Family: EUCHAETIDAE

- Genus; Euchaeta Philippi Euchaeta concinna Dana E. marina (Prestandrea) E. tenuis Esterly
 - E. wolfendeni A. Scott
- TRIBE: HETERARTHRANDRIA

Family: CENTROPAGIDAE

Genus: Centropages Kroyer Centropages furcatus (Dana) C. dorsispinatus Thompson and Scott C. alcocki Sewell

Family: PSEUDODIAPTOMIDAE

Genus: Pseudodiaptomus Herrick Pseudodiaptomus aurivilli Cleve

Genus: Schmackeria Poppe and Richard Schmackeria serricaudatus (T. Scott)

Family: TEMORIDAE

- Genus: Temora Baird Temora turbinata (Dana) T. discaudata Giesbrecht
- Family: METRIDIIDAE
- Genus: Pleuromamma Giesbrecht Pleuromamma indica Wolfenden
- Family: LUCICUTIIDAE
- Genus: Lucicutia Giesbrecht Lucicutia flavicornis (Claus)
- Family: ARIETELLIDAE
- Genus: Metacalanus Cleve Metacalanus aurivilli Cleve

Family: CANDACIIDAE

Genus: Candacia Dana Candacia bradyi A. Scott C. pachydactyla (Dana) Family: PONTELLIDAE

Genus: Labidocera Lubbock Labidocera acuta Dana L. pectinata Thompson and Scott L. minuta Giesbrecht

- L. pavo Giesbrecht
- Genus: Pontella Dana Pontella securifer Brady P. karachiensis Fazal-ur-Rehman P. spinipes Giesbrecht
- Genus: Pontellopsis Brady Pontellopsis herdmani Thompson and Scott P. macronyx A. Scott P. scotti Sewell

Genus: Pontellina Dana Pontellina plumata Dana Genus: Calanopia Dana Calanopia thompsoni A. Scott C. aurivilli (Cleve) C. minor A. Scott

Family: ACARTIIDAE

Genus: Acartia Dana

Acartia (Odontacartia) spinicaudata Giesbrecht A. (Odontacartia) centrura Giesbrecht A. (Odontacartia) erythraea Giesbrecht

- A. (Euacartia) southwelli Sewell

Family: TORTANIDAE

Genus: Tortanus Giesbrecht Tortanus gracilis (Brady T. forcipatus (Giesbrecht)

SUB ORDER: HARPACTICOIDA

SECTION: ACHIROTA

Family: ECTINOSOMIDAE

Genus: Microsetella Brady and Robertson Microsetella norvegica (Boeck)

Family: LONGIPEDIIDAE

Genus: Longipedia Claus Longipedia weberi A. Scott

SECTION: CHIROGNATHA

Family: TACHIDIIDAE

Genus: Euterpina Norman Euterpina acutifrons (Dana) Family: MACROSETELLIDAE

Genus: Macrosetella A. Scott Macrosetella gracilis (Dana)

Genus: Miracia Dana Miracia efferata Dana

Family: CLYTEMNESTRIDAE

Genus: Clyteninestra Dana Clytemnestra scutellata Dana

Family: METIDAE

Genus: Metis Philippi Metis jousseaumei (Richard)

SUR ORDER: CYCLOPOIDA

SECTION: GNATHOSTOMA

Family: OITHONIDAE

Genus: Oithona Baird Oithona brevicornis Giesbrecht O. rigida Giesbrecht O. similis Claus O. plumifera Baird O. spinirostris Claus

SECTION: POECILOSTOMA

Family: ONCAEIDAE

Genus: Oncaea Philippi Oncaea venusta Philippi O. conifera Giesbrecht

Family: SAPPHIRINIDAE

Genus: Sapphirina Thompson Sapphirina auronitens Claus S. ovatolanceolata Dana

Genus: Copilia Dana Copilia quadrata Dana C. mirabilis Dana

Family: CORYCAEIDAE

Genus: Corycaeus Dana

- Corycaeus (Corycaeus) crassiusculus Dana C. (Corycaeus) speciosus Dana C. (Ditrichocorycaeus) africanus F. Dahl C. (Ditrichocorycaeus) asiaticus F. Dahl C. (Onychocorycaeus) catus F. Dahl C. (Corycella) gibbulus Giesbrecht

660

Pontella karachiensis Fazal-ur-Rehman (Fig. 1)

Material : 1 female, 1 male, station 2, Off Port Okha, 6 m, 2 females, 1 male, station 3, Off Dwarka, 38m.




Fig. 1. Pontella karachiensis Fazal-ur-Rehman: a. lateral view (Female); b. right 1st antenna (Female); c. fifth leg (Female); d. lateral view (Male); e. right 1st antenna (Male); f. left fifth leg (Male); and g. right fifth leg (Male).

Description : Female : Total length 5.03 mm; length ratio of cephalothorax and abdomen is 62:39, so that abdomen is contained 1.564 times in the length of amerior region of body. Cephalon and 1st thoracic segment separate, as also 4th and 5th thoracic segments. Posterior thoracic segment is slightly flexed posteriorly, M. I. PATEL

nearly rounded and provided with a minute spine on each side. Abdomen asymmetrical and consists of two segments. Proportional lengths of segments are as follows:

Abdominal segment :
$$1-4$$
 5 Furca
 27 22 51 = 100

First antenna reaches back to about posterior margin of 2nd thoracic segment. It consists of 24 free segments having the following proportional lengths:

Segment	1	2	3	4	5	6	7	8	9	10	11	12
	36	53	18	2 8	18	18	24	21	24	28	21	21
Segment	13	14	15	16	17	18	19	2 0	2 1	22	23	24
	45	42	45	73	66	91	66	49	56	56	59	42 = 1000

Second antenna and mandible are as described and figured by Fazal-ur-Rehman (1973). Maxillules, maxillae and maxillipedes resemble those of other members of the genus. The swimming legs in the present specimens agree exactly with the figures given by Fazal-ur-Rehman (1973).

Male: Total length 4.08 mm. Body cylindrical. Proportional length of cephalothorax and abdomen 55:34, so that abdomen is contained 1.617 times in length of anterior region of body. Cephalic region, thoracic somites and other structures are similar in shape to that of female.

Abdomen symmetrical and consists of 5 segments. Proportional lengths of segments are as follows:

Abdominal segments:
$$1 \ 2 \ 3 \ 4 \ 5 \ Furca$$

 $18 \ 15 \ 19 \ 12 \ 10 \ 26 = 100$

First antenna on left side unmodified and consists usual 24 free joints. The flexion of geniculate right antenna situated between 18th and 19th segment. 17th segment bears an elongated ridge, with annulations proximally and fine denticles distally, extending to segment 16. 18th segment is armed with a similar but smaller ridge, carrying a single row of large spiniform denticles. Segmentation of 20th segment indistinct.

Details of fifth leg agrees exactly with original description of the species.

Distribution : Manora Channel, Karachi (Fazal-ur-Rehman, 1973).

REFERENCES

BHASKARAN, M. AND P. GOPALAKRISHNAN 1971. Observations on the marine plankton in the Gulf of Kutch, off Port Okha. Indian. J. Fish., 18: 99-108.

662

FAZAL-UR-REHMAN 1973. A new calanoid copepod Pontella karachiensis sp. nov. from West Pakistan. Crustaceana, 24 (2): 151-156.

PRASAD, R. R. 1964. A bibliography of the plankton of the Indian Ocean. INCOR (CSIR), New Delhi, Publication 3: 87 pp.

RAMAMURTHY, S. AND R. M. DHAWAN 1963. On the characteristics of the plankton at Kandla in the Gulf of Kutch during August 1958 to July 1960. Indian J. Fish., 10 (1): 94-101.

*SARS, G. O. 1903. Copepoda Calanoida. An Account of the Crustacea of Norway, 4: 1-171.

.

*_____ i911. Copepoda Harpacticoida. Ibid., 5: 1-449.

*_____ 1918. Copepoda Cyclopoida. Ibid., 6: 1-225.

*Not seen original reference.